

5% Sodium Fluoride Varnish

— Your partner in cavity prevention



Summary of advantages

• Stable and long-lasting fluorine release —

22,600 ppm fluoride

• Excellent consistency and long-term performance stability

• Fine and uniform particles enable a stable suspension of sodium fluoride, enhances fluorine release, as well as distribution

• Pleasant banana flavor to improve patients experience

• Unique packaging design facilitates clinical operation

Light yellow provides visual control
Material color: Light Yellow

Pro Shield Fluor Protector

To keep teeth healthy, small amounts of fluoride are needed during all stages of tooth growth. Several dental organizations and research reports recommend the professional use of fluoride varnish. Pro Shield is used for the prevention of recurrent cavities, cavity development, decalcification around orthodontic appliances, as well as pit and fissure (occlusal) cavities. Based on our patented technology, we come up with a

unique packaging. The design of separating the medicament and the brush head maximizes the doctor's operational convenience.







VINCISMILE GROUP LLC

Add:20524 Carrey Rd. Walnut, CA 91789, USA Email: marketing@vincismile.com Shop with us online at store.vincismile.com Tel: +1 626 283 5808 Web: www.vincismile.com



f 💿 D in Follow @VinciSmile

Excellent Operation Experience with Pro Shield



1. Make sure the areas to be treated are clean.



2. Remove excess saliva with triple syringe or cotton pellets.



3. For 10g tube: Squeeze an appropriate amount of material from the tube.



4. For 0.4g single dose: Apply the single dose directly.



5. Use the microbrush to apply a thin layer to the tooth surface.


2. Frequency of use:

Use once every 3-6 months.



6. Pro Sheild forms a thin film on the tooth surface.

High Fluoride Release Capability Suitable for Kids and Adults



Fluoride Release Rate (Long Term) 40.00% 35.00% 30.00% 25.00% 20.00% 15 00% 10.00% 5.00% 0.00% 10 15 20 25 30 0 time/h --- Pro Shield ----- B Varnish 🛶 A Varnish

*Data from VinciSmile Lab

Clinical application suggestions:

1. Recommended dosage:

Deciduous dentition <0.25g (5.65mg fluoride) Mixed dentition <0.4g (9.04mg fluoride) Permanent dentition <0.75g (16.95mg fluoride)

3. Precautions after use:

After using this product, do not brush the teeth, chew food, or drink hot drinks and alcoholic products (beverages, mouthwash, etc.) within **4 hours.**

